
#### The 2006 Boreal Forest Fire Season as seen by OMI

Omar Torres, Ray Hoff, Ray Rogers University of Maryland, Baltimore County

Changwoo Ahn, Colin Seftor SSAI, Lanham, Md

Aura Science Meeting Aerosols/Cloud/SO2 group 9-12-2006



OMI Aerosol Index 09-06-06

#### **OMI Near UV Aerosol Algorithm (OMAERUV)**

#### **Products:**

- -Near UV Aerosol Index
- -Extinction Optical Depth, AOD (388 nm)
- -Absorption Optical Depth, AAOD (388 nm)
- -AOD and AAOD reported also at 354 and 500 nm


#### Two sets of AOD and AAOD are reported

**Set 1**: As a function of assumed aerosol layer height above surface: 0.0, 1.5, 3.0. 6.0 and 10.0 km

**Set 2**: Recommended values for a particular assumption on aerosol layer height.

For boreal forest fires the aerosol layer is assumed to be at 6.0 km


# Absorption Optical Depth (388 nm), July 3-8, 2006





#### **OMI and CALIPSO Observations on July 5 and 7**



## Credits: CALIPSO Science Team










Credits: CALIPSO Science Team CALIPSO




July 10











## Arrival to the East Coast (July 7 and 8)




(10 km retrievals)


## Smoke vertical distribution over UMBC




### Credits Ray Rogers/UMBC





NOAA HYSPLIT MODEL Forward trajectories starting at 12 UTC 05 Jul 06 GDAS Meteorological Data




#### The Canadian smoke layer as detected from the ground










OMI Retrieved AOD (388 nm)

Forests Fires in Siberia July 23-30, 2006



#### Aeronet measurements on the path of the smoke layer

